

Flickor

Ålder	iso-bmi 25 ₁	iso-bmi 30 ₁	iso-bmi 35 ₂
2	18,02	19,81	22,9
3	17,56	19,36	22,3
4	17,28	19,15	22,0
5	17,15	19,17	22,3
6	17,34	19,65	23,2
7	17,75	20,51	24,6
8	18,35	21,57	26,5
9	19,07	22,81	28,8
10	19,86	24,11	31,2
11	20,74	25,42	33,4
12	21,68	26,67	34,9
13	22,58	27,76	35,9
14	23,34	28,57	36,3
15	23,94	29,11	36,4
16	24,37	29,43	36,1
17	24,70	29,69	35,7
18	25	30	35,0

Pojkar

Ålder	iso-bmi 25 ₁	iso-bmi 30 ₁	iso-bmi 35 ₂
2	18,41	20,09	22,5
3	17,89	19,57	21,7
4	17,55	19,29	21,3
5	17,42	19,30	21,3
6	17,55	19,78	21,6
7	17,92	20,63	22,3
8	18,44	21,60	23,5
9	19,10	22,77	25,1
10	19,84	24,00	27,2
11	20,55	25,10	29,7
12	21,22	26,02	32,3
13	21,91	26,84	34,6
14	22,62	27,63	36,1
15	23,29	28,30	36,8
16	23,90	28,88	36,6
17	24,46	29,41	35,9
18	25	30	35,0

Ref. ¹ Cole TJ et al BMJ vol 320, 1-6, 2000. (BMJ 320:1240-1243, 2000). ² Forslund et al.